arXiv:1606.01998v1 [math.CV] 7 Jun 2016

A REMARK ON EQUISINGULAR APPROXIMATIONS OF QUASI-PLURISUBHARMONIC FUNCTIONS

QI'AN GUAN AND ZHENQIAN LI

Abstract. In this note, we will present global equisingular approximations of quasi-plurisubharmonic functions with stable analytic pluripolar sets on compact complex manifolds.

1. Introduction

Let X be a compact complex manifold of dimension n. A function φ on X is said to be *quasi-plurisubharmonic (quasi-psh)* if φ is locally equal to the sum of a plurisubharmonic function and of a smooth function, that is, if its complex Hessian is bounded below by a $(1,1)$-form with continuous or locally bounded coefficients. The *multiplier ideal sheaf* $\mathscr{I}(\varphi)$ associated to a quasi-psh function φ is defined to be the sheaf of germs of holomorphic functions f such that $|f|^2e^{-2\varphi}$ is locally integrable with respect to the Lebesgue measure, which is a coherent ideal sheaf.

We say that a quasi-psh function φ has *analytic singularities* if φ can be written locally as

$$\varphi=\frac{c}{2}\log\big(\sum_{k=1}^{m}|f_k|^2\big)+O(1),$$

where $c\in\mathbb{R}^+$ and f_k are holomorphic functions.

In [1], Demailly proved a basic result on the approximation of psh functions by psh functions with analytic singularities via Bergman kernels. Here, we recall a global version as follows:

Theorem 1.1. ([1]; see also [2, 3]). *Let φ be a quasi-psh function on a compact Hermitian manifold (X,ω) of dimension n such that $\frac{i}{\pi}\partial\bar{\partial}\varphi\geq\gamma$ for some continuous $(1,1)$-form γ. Then there is a sequence of quasi-psh functions φ_m such that φ_m has the same singularities as a logarithm of a sum of squares of holomorphic functions and a decreasing sequence $\varepsilon_m>0$ converging to 0 such that*

(a) $\varphi(x)\leq\varphi_m(x)\leq\sup\limits_{|\zeta-x|<r}\varphi(\zeta)+C\big(\frac{|\log r|}{m}+r+\varepsilon_m\big)$ with respect to coordinate open sets covering X. In particular, φ_m converges to φ and in $L^1(X)$ and

(b) $\nu(\varphi,x)-\frac{n}{m}\leq\nu(\varphi_m,x)\leq\nu(\varphi,x)$ for every $x\in X$, where $\nu(\varphi,x)$ is the Lelong number of φ at x;

(c) $\frac{i}{\pi}\partial\bar{\partial}\varphi\geq\gamma-\varepsilon_m\omega$.

In this note, by the strong openness property of multiplier ideal sheaves [6], we will present the following equisingular approximations quasi-psh functions with

Date: June 8, 2016.

2010 *Mathematics Subject Classification.* 32C25, 32C35, 32U05.

The first author was partially supported by NSFC-11522101 and NSFC-11431013.

Key words. Plurisubharmonic function, Multiplier ideal sheaf, Equisingularity.

stable analytic pluripolar set $A := N(\mathscr{I}(\varphi))$, the zero set of multiplier ideal sheaf $\mathscr{I}(\varphi)$ associated to φ.

Theorem 1.2. *Let φ be a quasi-psh function on a compact complex manifold X. Then, there exists a sequence $(\hat{\psi}_k)$ of quasi-psh functions with analytic singularities on X such that*

(1) *$(\hat{\psi}_k)$ converges to φ almost everywhere;*
(2) ("*equisingularity*") *$\mathscr{I}(\hat{\psi}_k) = \mathscr{I}(\varphi)$ for every k;*
(3) ("*stability*") *the polar sets of $\hat{\psi}_k$ coincide with A for all k.*

Remark 1.1. In [5], the first author proved that one cannot add the requirement to make $\hat{\psi}_k$ converge to φ decreasingly.

In [4], the following "equisingular" regularization process was established, which provides more equisingularity than Theorem 1.1 in the sense that the multiplier ideal sheaves are preserved.

Theorem 1.3. ([4], see also [2, 3]). *Let $T = \alpha + i\partial\bar{\partial}\varphi$ be a closed $(1,1)$-current on a compact Hermitian manifold (X,ω), where α is a smooth closed $(1,1)$-form and φ a quasi-psh function. Let γ be a continuous real $(1,1)$-form such that $T \geq \gamma$. Then one can write $\varphi = \lim_{m\to+\infty}\varphi_m$, where*

(a) *φ_m is smooth in the complement $X\backslash Z_m$ of an analytic set $Z_m \subset X$;*
(b) *(φ_m) is a decreasing sequence, and $Z_m \subset Z_{m+1}$ for all m;*
(c) *$\int_X (e^{-2\varphi} - e^{-2\varphi_m})dV_\omega$ is finite for every m and converges to 0 as $m \to +\infty$;*
(d) ("*equisingularity*") *$\mathscr{I}(\varphi_m) = \mathscr{I}(\varphi)$ for all m;*
(e) *$T_m = \alpha + i\partial\bar{\partial}\varphi_m$ satisfies $T_m \geq \gamma - \varepsilon_m\omega$, where $\varepsilon_m \to 0$ as $m \to +\infty$.*

Note that in Theorem 1.3 the pluripolar set Z_m of φ_m is larger and larger as m tends to infinity. Combining the strong openness property of multiplier ideal sheaves and Theorem 1.3, we also present the following

Theorem 1.4. *Let φ be a quasi-psh function on a compact complex manifold X. Then, there exists a sequence $(\hat{\psi}_k)$ of quasi-psh functions on X such that*

(1) *$(\hat{\psi}_k)$ is decreasing and convergent to φ;*
(2) *$\hat{\psi}_k$ is smooth on $X\backslash A$ for all k;*
(3) *$e^{-2\varphi} - e^{-2\hat{\psi}_k}$ is locally integrable for every k, which implies $\mathscr{I}(\hat{\psi}_k) = \mathscr{I}(\varphi)$.*

2. Proof of main results

To prove main results, the following strong openness property of multiplier ideal sheaves is necessary.

Theorem 2.1. ([6]). *Let φ be a quasi-psh function on complex manifold X and $K \subset X$ a compact subset. Then, there exists $\varepsilon_K > 0$ such that for any $0 \leq \varepsilon \leq \varepsilon_K$ we have*

$$\mathscr{I}((1+\varepsilon)\varphi)|_K = \mathscr{I}(\varphi)|_K.$$

We are now in a position to prove our main results.

***Proof of Theorem* 1.2.** Step 1. By Theorem 1.1, there exists a finite open covering (U_k) of X by coordinate balls and a sequence of quasi-psh functions φ_m

on X such that on every U_k, we have

$$\varphi_m|_{U_k} = \frac{1}{2m}\log\sum_j |\sigma_{j,m}^{(k)}|^2 + O(1),$$

where $(\sigma_{j,m}^{(k)})$ is an orthonormal basis of $\mathcal{H}_{U_k}(m\varphi)$, the Hilbert space of holomorphic functions f on U_k satisfying $\int_{U_k}|f|^2e^{-2m\varphi}d\lambda_n<\infty$.

Step 2 (see Proposition 2.1 in [7]). There exists a quasi-psh function $\tilde{\varphi}$ on X such that

(i) $e^{-2\varphi}-e^{-2\tilde{\varphi}}$ is locally integrable, which implies $\mathscr{I}(\varphi)=\mathscr{I}(\tilde{\varphi})$ on X;

(ii) $\tilde{\varphi}\in L^\infty_{loc}(X\backslash A)$.

Let $x_0\in X$ be any point. Without loss of generality, we assume $x_0\in U_k$. By Proposition 5.7 in [2], there exists $j_0>0$ and a neighborhood $V_0\subset\subset U_k$ of x_0 such that $\sigma_{1,1}^{(k)},...,\sigma_{j_0,1}^{(k)}$ generate $\mathscr{I}(\varphi)$ on V_0, and

$$\log\sum_j|\sigma_{j,1}^{(k)}|^2=\log\sum_{j=1}^{j_0}|\sigma_{j,1}^{(k)}|^2+O(1).$$

on U_k. It follows from Theorem 2.1 that there exists a real number $p_0>0$ satisfying

$$\int_{V_0}|\sigma_{j,1}^{(k)}|^2e^{-2(1+\frac{1}{p})\varphi}d\lambda_n<\infty$$

for any $1\leq j\leq j_0$ and any $p\geq p_0$. Then, we obtain that on V_0,

$$\begin{aligned}&\int_{V_0}(e^{-2\varphi}-e^{-2\max\{\varphi,p\varphi_1\}})d\lambda_n=\int_{\{\varphi<p\varphi_1\}\cap V_0}e^{2\cdot\frac{1}{p}\varphi-2(1+\frac{1}{p})\varphi}d\lambda_n\\ \leq&\int_{\{\varphi<p\varphi_1\}\cap V_0}e^{2\varphi_1-2(1+\frac{1}{p})\varphi}d\lambda_n\leq\int_{V_0}e^{2\varphi_1-2(1+\frac{1}{p})\varphi}d\lambda_n\\ \leq&C\cdot\int_{V_0}\sum_{j=1}^{j_0}|\sigma_{j,m}^{(k)}|^2e^{-2(1+\frac{1}{p})\varphi}d\lambda_n<\infty\end{aligned}$$

for some constant $C>0$.

Thus, we infer from the compactness of X that for sufficiently large p_0, by taking $\tilde{\varphi}:=\max\{\varphi,p\varphi_1\}$ with $p\geq p_0$, we have $e^{-2\varphi}-e^{-2\tilde{\varphi}}$ is locally integrable and $\tilde{\varphi}\in L^\infty_{loc}(X\backslash A)$.

Step 3. As the discussion of Remark 3 in [3], by taking $\hat{\varphi}_m=(1+\frac{1}{m})\tilde{\varphi}_m$, we have $\mathscr{I}(\hat{\varphi}_m)=\mathscr{I}(\tilde{\varphi})$ for large enough m, where $\tilde{\varphi}_m$ is the Bergman approximation sequence of $\tilde{\varphi}$. Moreover, $\hat{\varphi}_m$ is smooth outside the polar set Z_m of $\hat{\varphi}_m$.

Since $\hat{\varphi}_m|_{X\backslash A}$ is smooth, we have $Z_m\subset A$. Then, it follows from

$$\mathscr{I}(\hat{\varphi}_m)_x=\mathscr{I}(\tilde{\varphi})_x\neq\mathcal{O}_{X,x}$$

for any $x\in A$ that $Z_m=A$ for large enough m.

Step 4. Equisingular approximation of φ.

Without loss of generality, we assume that both φ and φ_1 are negative on X. Let $\psi_k:=\max\{\varphi,p_k\varphi_1\}$, where $p_0\leq p_k<p_{k+1}\to\infty$ $(k\to\infty)$. Then ψ_k converges

to φ and $\mathscr{I}(\psi_k) = \mathscr{I}(\varphi)$ for every k. By Remark 3 in [3], for every k, there exists a sequence of quasi-psh functions ψ_{jk} with analytic singularities such that $\mathscr{I}(\psi_{jk}) = \mathscr{I}(\varphi_k)$ for every j, and ψ_{jk} is convergent to φ_k. It follows that for every k, there exists $N_k > 0$ such that $m(E\{|\psi_{jk} - \psi_k| \geq \frac{1}{2^k}\}) \leq \frac{1}{2^k}$, for any $j \geq N_k$.

Take a subsequence $\psi_{j_k k}$ of ψ_{jk} with $N_k \leq j_k < j_{k+1}$. It follows from

$$|\psi_{j_k k} - \varphi| \leq |\psi_{j_k k} - \psi_k| + |\psi_k - \varphi|$$

that for every $\varepsilon > 0$ and all k with $\frac{1}{2^k} < \varepsilon$, we have

$$E\{|\psi_{j_k k} - \varphi| \geq \varepsilon\} \subset E\{|\psi_{j_k k} - \psi_k| \geq \frac{1}{2^k}\} \cup E\{|\psi_k - \varphi| \geq \varepsilon - \frac{1}{2^k}\},$$

which implies

$$m(E\{|\psi_{j_k k} - \varphi| \geq \varepsilon\}) \leq \frac{1}{2^k} + \frac{\varepsilon}{2} < \varepsilon$$

for large enough k, i.e., $\psi_{j_k k}$ is convergent to φ in Lebesgue measure. Let $\hat{\psi}_k$ be a subsequence of $\psi_{j_k k}$, which converges to φ almost everywhere. Then, it follows from Step 3 that $\hat{\psi}_k$ satisfies (1), (2) and (3) as desired. □

***Proof of Theorem* 1.4.** Let (ψ_k) be the equisingular approximation sequence of φ as in Theorem 1.3 and φ_1, p_0 as in Step 2 of Theorem 1.2. In addition, we can assume all ψ_k and φ_1 are negative. Then, we obtain that $e^{-2\psi_k} - e^{-2\varphi}$ and $e^{-2\varphi} - e^{-2\max\{\varphi, p\varphi_1\}}$ are locally integrable for any $p \geq p_0$, which implies that $e^{-2\psi_k} - e^{-2\max\{\varphi, p\varphi_1\}}$ is locally integrable. Since

$$e^{-2\psi_k} - e^{-2\max\{\psi_k, \max\{\varphi, p\varphi_1\}\}} = \max\{0, e^{-2\psi_k} - e^{-2\max\{\varphi, p\varphi_1\}}\} \in L^1_{loc}(X),$$

then $e^{-2\psi_k} - e^{-2\max\{\psi_k, p\varphi_1\}}$ is locally integrable.

Let

$$M_\eta(t_1, t_2) := \int_{\mathbb{R}^2} \max\{t_1 + x_1, t_2 + x_2\} \prod_{1 \leq j \leq 2} \eta_j^{-1} \theta(x_j/\eta_j) dx_1 dx_2$$

be the regularized max function, where $\eta = (\eta_1, \eta_2)$ with $\eta_j > 0$, and θ is a non-negative smooth function on $\mathbb{R}$ with support in $[-1, 1]$ such that $\int_{\mathbb{R}} \theta(x) dx = 1$ and $\int_{\mathbb{R}} x\theta(x) dx = 0$. By setting $\eta = (1, 1)$, we have $M_\eta(\max\{\psi_k, p_0\varphi_1\}, p\varphi_1) = \max\{\psi_k, p_0\varphi_1\}$ near A for any $p > p_0$. Then, it follows that

$$e^{-2\max\{\psi_k, p_0\varphi_1\}} - e^{-2M_\eta(\max\{\psi_k, p_0\varphi_1\}, p\varphi_1)}$$

is locally integrable.

Note that

$$\begin{aligned} 0 \leq & e^{-2\psi_k} - e^{-2M_\eta(\psi_k, p\varphi_1)} \\ \leq & e^{-2\psi_k} - e^{-2M_\eta(\max\{\psi_k, p_0\varphi_1\}, p\varphi_1)} \\ = & (e^{-2\psi_k} - e^{-2\max\{\psi_k, p_0\varphi_1\}}) + (e^{-2\max\{\psi_k, p_0\varphi_1\}} - e^{-2M_\eta(\max\{\psi_k, p_0\varphi_1\}, p\varphi_1)}) \end{aligned}$$

for any $p > p_0$. Thus, $e^{-2\psi_k} - e^{-2M_\eta(\psi_k, p\varphi_1)}$ is locally integrable for any $p > p_0$ and all k.

Let $\hat{\psi}_k := M_\eta(\psi_k, p_k\varphi_1)$, where $\eta = (1, 1)$ and $p_0 < p_k < p_{k+1} \to \infty$ $(k \to \infty)$. Then, $\hat{\psi}_k$ decreasingly converges to φ and smooth on $X \backslash A$. It follows from

$$e^{-2\varphi} - e^{-2\hat{\psi}_k} = (e^{-2\varphi} - e^{-2\psi_k}) + (e^{-2\psi_k} - e^{-2M_\eta(\psi_k, p_k\varphi_1)})$$

that $e^{-2\varphi} - e^{-2\hat{\psi}_k}$ is locally integrable. □

References

[1] J.-P. Demailly, *Regularization of closed positive currents and intersection theory*, J. Algebraic Geom. 1 (1992), 361–409.

[2] J.-P. Demailly, *Analytic Methods in Algebraic Geometry*, Higher Education Press, Beijing, 2010.

[3] J.-P. Demailly, *On the Cohomology of Pseudoeffective Line Bundles*, Complex Geometry and Dynamics, The Abel Symposium 2013, 51–99.

[4] J.-P. Demailly, T. Peternell, M. Schneider, *Pseudo-effective line bundles on compact Kähler manifolds*, Internat. J. Math. 12 (2001), 689–741.

[5] Q. A. Guan, *Nonexistence of decreasing equisingular approximations with logarithmic poles*, J. Geom. Anal., published online on March 16, 2016; DOI: 10.1007/s12220-016-9702-2.

[6] Q. A. Guan, X. Y. Zhou, *A proof of Demailly's strong openness conjecture*, Ann. of Math. (2) 182 (2015), 605–616. See also arXiv:1311.3781.

[7] Q. A. Guan, X. Y. Zhou, *Characterization of multiplier ideal sheaves with weights of Lelong number one*, Adv. Math. 285 (2015), 1688–1705.

Qi'an Guan: School of Mathematical Sciences, and Beijing International Center for Mathematical Research, Peking University, Beijing, 100871, China.

E-mail address: guanqian@amss.ac.cn

Zhenqian Li: School of Mathematical Sciences, Peking University, Beijing, 100871, China.

E-mail address: lizhenqian@amss.ac.cn